

IMMUNIZATIONS

Vaccinations begin soon after birth and, for the most part, are completed by age two. It is important to catch up if an immunization is missed. Don't just skip it. The schedule shown below is provided by the Center for Disease Control (CDC). The administering physician will determine if any change in vaccine is necessary on a case-by-case basis.

Vaccine	Hepatitis B	Diphtheria Tetanus Pertussis	Haemophilus Influenzae type b	Inactivated Poliovirus	Measles Mumps Rubella	Varicella (Chicken Pox)	Pneumo- coccal Conjugate	Influenza
Birth								
1 mo.								
2 mo.								
4 mo.								
6 mo.								
12 mo.								
15 mo.								
18 mo.								
24 mo.								
4-6 yr.								
11-12 yr.								
13-18 yr.								

For a current recommended childhood immunization schedule from the Center for Disease Control, go to www.cdc.gov/nip on the web.

RECOMMENDED ADULT IMMUNIZATION SCHEDULE

	19-49	50-64	Over 65
Tetanus, Diphtheria	1 dose booster every 10 years for all persons over age 19		
Influenza	1 dose annually if at risk	1 dose annually	
Pneumococcal	1 dose for all persons at risk		1 dose
Hepatitis B	3 doses for all persons at risk over age 19		
Hepatitis A	2 doses for all persons at risk over age 19		
Measles, Mumps, Rubella	1 or 2 doses (if no previous vaccination or has symptoms)		
Varicella	2 doses if no previous vaccination or has symptoms		
Meningococcal	1 dose for persons at risk over age 19		

